

Neutrino flavor conversion in a neutrino background: single- versus multi-particle description

Alexander Friedland^{1,2,*} and Cecilia Lunardini^{2,†}

¹Theoretical Division, T-8, MS B285, Los Alamos National Laboratory, Los Alamos, NM 87545

²School of Natural Sciences, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540

(Dated: February 1, 2008)

In the early Universe, or near a supernova core, neutrino flavor evolution may be affected by coherent neutrino-neutrino scattering. We develop a microscopic picture of this phenomenon. We show that coherent scattering does not lead to the formation of entangled states in the neutrino ensemble and therefore the evolution of the system can always be described by a set of one-particle equations. We also show that the previously accepted formalism overcounts the neutrino interaction energy; the correct one-particle evolution equations for both active-active and active-sterile oscillations contain additional terms. These additional terms modify the index of refraction of the neutrino medium, but have no effect on oscillation physics.

PACS numbers: 14.60.Pq

I. INTRODUCTION

It is well known that interaction with a medium modifies neutrino dispersion relations or, equivalently, gives the medium a nontrivial refraction index for neutrinos. Because this effect is generically flavor-dependent, it can have a profound impact on neutrino flavor evolution. Inside the Sun and the Earth, the refraction effect arises as a result of neutrino interactions with electrons and nucleons. On the other hand, in the early Universe or near a supernova core, where the number density of neutrinos is sufficiently high, the refractive properties of the neutrino background itself are important [1, 2, 3, 4, 5]. This neutrino “self-refraction” is the subject of the present work.

Early studies [2, 3, 4, 5, 6, 7] of the effect treated the neutrino background analogously to the case of ordinary matter (electrons and nucleons), namely, assuming that

- for each neutrino, one can write a single-particle evolution equation and take into account the effect of all other particles (including other neutrinos) by adding appropriate terms to the one-particle Hamiltonian;
- the contribution of the background neutrinos to this Hamiltonian is diagonal in the flavor basis.

With these assumptions, the evolution of each neutrino in the system was described by

$$i \frac{d\nu^{(i)}}{dt} = (H_{\text{vac}} + H_{\text{mat}} + \sum_j H_{\nu\nu}^{(ij)})\nu^{(i)}, \quad (1)$$

where i and j label neutrinos in the system, H_{vac} and H_{mat} are the usual vacuum and ordinary matter Hamiltonian terms and the last term is the sum over the contri-

butions of all background neutrinos, taken to be flavor-diagonal.

These assumptions were critically reexamined by James Pantaleone [8, 9] who made several important observations regarding their validity. First, he reasoned that flavor evolution of neutrinos in a neutrino background is in general a many-body phenomenon and it is *a priori* not obvious that the first assumption is justified in all cases. Second, he demonstrated that, for a system of several active neutrinos flavors, even when the first assumption is justified, the second one is incompatible with the symmetry of the problem. Indeed, the low energy neutral current Hamiltonian [30],

$$H_{\text{NC}} = \frac{G_F}{\sqrt{2}} \left(\sum_a j_a^\mu \right) \left(\sum_b j_{b\mu} \right), \quad (2)$$

where the currents $j_a^\mu \equiv \bar{\nu}_a \gamma^\mu \nu_a$ and a is a flavor index, $a = 1, \dots, N$, possesses a $U(N)$ flavor symmetry, which must be respected by any effective description of the system. This requirement is not satisfied by the diagonal form for $H_{\nu\nu}^{(ij)}$ used in the earlier studies.

Pantaleone proposed a modified form for $H_{\nu\nu}^{(ij)}$, which contained non-zero off-diagonal terms,

$$H_{\nu\nu}^{(ij)} = A \left[|\nu_e^{(j)}|^2 + |\nu_\mu^{(j)}|^2 + \begin{pmatrix} |\nu_e^{(j)}|^2 & \nu_e^{(j)} \nu_\mu^{(j)*} \\ \nu_e^{(j)*} \nu_\mu^{(j)} & |\nu_\mu^{(j)}|^2 \end{pmatrix} \right]. \quad (3)$$

In Eq. (3) the wave function of the background neutrino $\nu^{(j)}$ is normalized such that $\int dV (|\nu_e^{(j)}|^2 + |\nu_\mu^{(j)}|^2) = 1$ and the coefficient of proportionality A equals $\sqrt{2}G_F(1 - \cos \beta^{(ij)})$, $\beta^{(ij)}$ being the angle between the two neutrino momenta. For antineutrinos in the background, the form of the Hamiltonian is exactly the same, with the only difference that A has the opposite sign. As can be easily checked, Eq. (3) is indeed consistent with the $U(2)$ flavor symmetry of the two-flavor system.

The result (3) was also later obtained by Sigl and Raffelt [10] and by McKellar and Thomson [11] in the frame-

*Electronic address: friedland@lanl.gov

†Electronic address: lunardi@ias.edu

work of more general analyses of the flavor evolution of a neutrino system, which took into account both refraction effects and collisions. Eqs. (1,3) have been used as a starting point for extensive studies of the neutrino evolution in the early Universe [12, 13, 14, 15, 16, 17, 18, 19, 20] and in supernovae [21, 22, 23, 24, 25]. The general properties of the solutions have been investigated in [26, 27, 28, 29].

Certain theoretical aspects of these equations, however, have not received adequate attention in the literature to date. One such aspect is the validity of a description of the problem by a set of single-particle equations. The absence of quantum correlations (entanglement) at all times during the evolution of the system is *a priori* not obvious, and has not been proven. Indeed, in the derivations [10] and [11] it is stated as an assumption. In [9], it is suggested that for quantum correlations not to form the system must obey certain physical conditions, for example, it should contain an incoherent mixture of mass eigenstates. This raises the question how general the results in Eqs. (1,3) are and what physical criteria determine their breakdown.

The second important aspect is the connection between the elementary neutrino-neutrino scattering processes and the macroscopic description of refraction given in Eqs. (1,3). For refraction in ordinary matter, such a connection is very well established and provides the most straightforward derivation of the effect as an interference of many elementary scattering amplitudes. A similar treatment for the case of the neutrino background has not been given.

In this paper we present a description of the flavor evolution of neutrinos in a neutrino background in terms of the elementary neutrino-neutrino scattering processes. This description provides a transparent, physical picture of the effect, and, at the same time, allows us to prove that quantum correlations in the neutrino ensemble are negligible and therefore the description in terms of one-particle equations is valid. These equations are obtained directly from our formalism and are compared to the accepted results, Eqs. (1,3).

We work in the regime in which (i) neutrino-neutrino interactions are described by the low-energy four-fermion Fermi coupling, (ii) the neutrino gas is non-degenerate, and (iii) incoherent scattering of neutrinos with other neutrinos and other particle species is negligible. The first condition implies that the neutrino center-of-mass energies are well below the weak scale. The second one is satisfied if a gas of neutrinos has a number density $n_\nu \ll E^3$, where E is a typical neutrino energy. Finally, incoherent scattering is significant if the column density of the medium exceeds the inverse of the scattering cross section: $d \equiv \int n(x)dx \gtrsim 1/\sigma$, as happens, e.g., in the early Universe at temperatures larger than few MeV (see, e.g., [18]).

The refraction effects of the neutrino background are negligible if the coupling between a neutrino and the neutrino gas is significantly smaller than the vacuum oscillation Hamiltonian or than the coupling to ordinary mat-




FIG. 1: Neutrino beam traversing a slab of FCNC interacting matter.

ter. This condition thus depends on the neutrino energy, on the density of the neutrino background, and on the density of the ordinary matter. In the core of the Sun, where the number density of neutrinos is $n_\nu \simeq 10^6 \text{ cm}^{-3}$, neutrino-neutrino interaction is negligible for all relevant neutrino energies and oscillation parameters. In contrast, near a supernova core the neutrino density can be as high as $n_\nu \simeq 10^{31} \text{ cm}^{-3}$ and is comparable with the electron density there. Therefore, neutrino-induced refraction effects are important and must be taken into account.

The paper is organized as follows. In Sect. II we review the microscopical picture of the neutrino flavor evolution in normal matter and point out that a naive extension of this picture to the neutrino self-refraction leads to seemingly paradoxical results. In Sect. III we show how this picture should be constructed consistently by identifying states in the neutrino ensemble that amplify coherently. We also show that coherent scattering does not form entangled states. In Sect. IV we compare the one-particle evolution equations we obtain for active-active and active-sterile oscillations in the neutrino and antineutrino backgrounds with the accepted results. In Sect. V we show how entanglement is effectively destroyed by the refraction phenomenon. Finally, Sect. VI summarizes our conclusions.

II. FORMULATION OF THE PROBLEM

A. Conventional neutrino refraction: FCNC case

We begin by briefly reviewing the physics of the refraction effect in “normal” matter. While there are many ways to derive the refraction properties in this case, fundamentally, the relevant effect is the coherent interference of many elementary scattering events.

Let us consider the problem of neutrino oscillations in a medium which possesses flavor changing neutral current (FCNC) interactions. Such interactions give rise to the nonzero off-diagonal terms in the neutrino evolution

Hamiltonian,

$$H_{FCNC} = \frac{\sqrt{2}G_F n_2}{2} \left[\text{const} + \begin{pmatrix} \epsilon' & \epsilon \\ \epsilon & -\epsilon' \end{pmatrix} \right], \quad (4)$$

where G_F is the Fermi constant and n_2 is the number density of scatterers in the medium.

As a toy example, consider a beam of electron neutrinos incident on a thin slab of matter of thickness L made of FCNC interacting particles, as illustrated in Fig. 1. Assume that the neutrino masses are sufficiently small so that the effects of vacuum oscillation can be neglected. The flavor conversion rate in the slab can then be found using the following straightforward physical argument. Let f be the amplitude for an electron neutrino to scatter as a muon neutrino in a given direction on a particle in the target. If the scattering amplitudes for different target particles add up incoherently, the flux of muon neutrinos in that direction is $\propto N_s |f|^2$, where N_s is the number of scatterers. In the case of forward scattering, however, the scattering amplitudes add up *coherently* and, hence, the forward flux of muon neutrinos is $\propto N_s^2 |f|^2$. Indeed, in the small L limit Eq. (4) gives

$$P_{\nu_e \rightarrow \nu_\mu}^{FCNC} \simeq \epsilon^2 (G_F n_2 L)^2 / 2, \quad (5)$$

which has the form $P_{\nu_e \rightarrow \nu_\mu}^{FCNC} \propto N_s^2 |f|^2$, since $\epsilon \propto f$. Notice that by choosing a small L limit we were able to ignore the secondary conversion effects in the slab, *i.e.*, to assume that for all elementary scattering events the incident neutrinos are in the ν_e state.

To summarize, for small enough L , the flavor conversion rate due to coherent FC scattering in the forward direction is proportional to the square of the modulus of the product of the elementary scattering amplitude and number of scatterers. This quadratic dependence on N_s is what makes the coherent forward scattering important even when the incoherent scattering can be neglected.

Notice that exactly the same arguments apply if one considers the usual flavor-diagonal matter term due to the electron background in a rotated basis, for instance, in the basis of vacuum mass eigenstates. In this basis, the matter Hamiltonian has off-diagonal terms, resulting in transitions between the vacuum mass eigenstates.

B. Neutrino background: physical introduction

We seek the same description for the case of neutrino background. Let us therefore modify the setup in Fig. 1 and replace the slab by a second neutrino beam, such that the neutrino momenta in the two beams are orthogonal (see Fig. 2). To keep the parallel between this case and the FCNC case, we will continue to refer to the original beam as “the beam” and to the second beam as “the background”. The neutrinos in each beam can be taken to be approximately monoenergetic [31]. We again assume that the neutrino masses are sufficiently small so




FIG. 2: Toy problem to illustrate neutrino flavor conversion in the neutrino background.




FIG. 3: Elementary scattering event that causes a change of the flavor composition of the beam

that, although flavor superposition states could be created *outside* the intersection region, the effects of vacuum oscillation *inside* the intersection region can be neglected. Any flavor conversion that takes place in the system is therefore due to neutrino-neutrino interactions in the intersection region.

Let us first compute the amount of flavor conversion in the beam using Eqs. (1,3). The conversion is expected because of the presence of the off-diagonal terms in these equations. The result depends on the flavor composition of the background. If the background neutrinos are all in the same flavor state

$$\nu_x = \cos \alpha \nu_e + \sin \alpha \nu_\mu \quad (6)$$

and their density is n_2 , the Hamiltonian for the evolution of a beam neutrino takes the form

$$H = \frac{\sqrt{2}G_F n_2}{2} \left[\text{const} + \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{pmatrix} \right]. \quad (7)$$

After traversing the intersection region, a neutrino in the beam will be converted to the ν_μ state with the probability

$$P_{\nu_e \rightarrow \nu_\mu} \simeq \sin^2 2\alpha (G_F n_2 L)^2 / 2, \quad (8)$$

assuming as before that the size of the region L is small.

The above descriptions of neutrino flavor conversion in the neutrino background and in the FCNC background are very similar in form. We will next show, however, that despite very similar appearances of the equations, the underlying physics in the two cases is different.

First, we need to establish which elementary processes give rise to neutrino flavor conversion in the neutrino background. In the absence of vacuum oscillations, the only interactions in the problem are neutral current inter-

actions between pairs of neutrinos. These are described by the Hamiltonian (2), which conserves the *total* flavor of the system. The flavor composition of the beam therefore changes only if some of the background neutrinos of different flavor scatter into the beam, *i.e.*, if a neutrino from the background and a neutrino from the beam *exchange* momenta, $\nu(\vec{k}) + \nu(\vec{p}) \rightarrow \nu(\vec{p}) + \nu(\vec{k})$. As observed in [9], such events can add up coherently and give rise to the flavor off-diagonal entries in the oscillation Hamiltonian.

Let us consider such an elementary event, as depicted in Fig. 3. Following [9], we restrict scattering angles to the directions in which coherent interference occurs and write the Hamiltonian for an interacting pair of neutrinos in the form

$$i \frac{d}{dt} \begin{pmatrix} |\nu_e(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_e(\vec{k})\nu_\mu(\vec{p})\rangle \\ |\nu_\mu(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_\mu(\vec{k})\nu_\mu(\vec{p})\rangle \end{pmatrix} = \frac{\sqrt{2}G_F}{V}(1 - \cos\beta) \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} |\nu_e(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_e(\vec{k})\nu_\mu(\vec{p})\rangle \\ |\nu_\mu(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_\mu(\vec{k})\nu_\mu(\vec{p})\rangle \end{pmatrix}, \quad (9)$$

where V is the normalization volume. For a state $|S\rangle$, which at $t = 0$ is $|\nu_e\nu_\mu\rangle$, this equation formally has a solution

$$|S(t)\rangle = \frac{|\nu_e\nu_\mu\rangle - |\nu_\mu\nu_e\rangle}{2} + e^{-i\delta Et} \frac{|\nu_e\nu_\mu\rangle + |\nu_\mu\nu_e\rangle}{2}, \quad (10)$$

where $\delta E = 2\sqrt{2}G_F(1 - \cos\beta)/V$. In practice, however, one only needs a small t expansion of Eq. (10):

$$\begin{aligned} S(t) &\simeq |\nu_e\nu_\mu\rangle - i\delta Et/2(|\nu_e\nu_\mu\rangle + |\nu_\mu\nu_e\rangle) \\ &= (1 + ia)|\nu_e\nu_\mu\rangle + ia|\nu_\mu\nu_e\rangle, \end{aligned} \quad (11)$$

where $a = -\sqrt{2}G_F(1 - \cos\beta)t/V$. Indeed, since the interaction time and the normalization volume are determined by the size of the neutrino wave packet l , $t \sim l$ and $V \sim l^3$, the absolute value of a in any realistic physical situation is always much less than 1. This simply means that the interaction between two neutrinos is described by the lowest order Feynman diagram and second order scattering effects may be safely ignored.

We now return to the situation depicted in Fig. 2. We are interested in the muon neutrino content of the beam after it crosses the interaction region. Using the Hamiltonian in Eq. (9) to describe an elementary scattering event $\nu_e + \nu_x$, we find

$$\begin{aligned} \tilde{S}(t) &\simeq |\nu_e\nu_x\rangle + ia(|\nu_x\nu_e\rangle + |\nu_e\nu_x\rangle), \\ &= (1 + ia)|\nu_e\nu_x\rangle + ia \cos\alpha |\nu_e\nu_e\rangle \\ &\quad + ia \sin\alpha |\nu_\mu\nu_e\rangle. \end{aligned} \quad (12)$$

Thus, for an elementary event, the amplitude of measuring the neutrino in the beam as ν_μ is proportional

to $\sin\alpha$. If one multiplies this amplitude by the number of scatterers and squares, as was previously done in the FCNC example, one finds $P_{\nu_e \rightarrow \nu_\mu} \propto N_s^2 \sin^2 \alpha$, which is in clear conflict with the prediction of Eq. (8), $P_{\nu_e \rightarrow \nu_\mu} \propto N_s^2 \sin^2 2\alpha$.

Which of the two results is right? At first sight, the first possibility may appear more plausible. Consider, for instance, what happens when an electron neutrino propagates through a background of muon neutrinos ($\alpha = \pi/2$). Since muon neutrinos appear in the beam as a result of elementary exchanges between the beam and the background, one may think that for a pure muon neutrino background the conversion rate in the beam should be maximal. The first result indeed has this behavior, while the second result predicts *no conversion*. One may therefore be tempted to conclude that the second result fails to describe the system in this limit and perhaps that this failure signals the general breakdown of the validity of the single-particle description.

As will be shown in the next section, however, despite its seemingly paradoxical behavior, the second result is actually correct. The explanation of the paradox lies in the procedure of adding up elementary amplitudes, which in the case of the neutrino background is more subtle than in the case of ordinary matter.

III. MICROSCOPIC ANALYSIS OF THE EFFECT

The key observation is that a neutrino-neutrino scattering event changes not only the state of the beam, but also the state of the background. This implies that the refraction by the neutrino background is intrinsically different from the case of ordinary matter, and requires a specific analysis to establish what states in the neutrino ensemble are amplified coherently.

We begin by recalling that an elementary scattering event can only be amplified if the particles scatter “forward”, meaning that either particle momenta do not change (t -channel), or that they are exchanged (u -channel) as in Fig. 3. The interaction is described by Eq. (11). For the sake of clarity, in this section we consider the exchange diagrams only and omit the effects of the non-exchange diagrams. The latter produce an overall (flavor-independent) phase shift of the neutrino states, and do not affect the neutrino flavor conversion.

We first consider a single electron neutrino in the beam interacting with several neutrinos in the background. Assume that these background neutrinos are all in the same flavor state ν_x , Eq. (6), and that the state of the background initially is a product of single-particle states, $|xxx...xx\rangle$. The total system therefore is initially in the state $|e\rangle|xxx...xx\rangle$ and, as a result of the interaction, evolves over a time step δt according to

$$|e\rangle|xxx...xx\rangle \longrightarrow |F\rangle = |e\rangle|xxx...xx\rangle + ia|F_1\rangle, \quad (13)$$

where $|F_1\rangle$ is the state with all possible exchange terms,

$$\begin{aligned} |F_1\rangle &= |x\rangle|exx...xx\rangle + |x\rangle|xex...xx\rangle \\ &+ |x\rangle|xex...xx\rangle + \dots \end{aligned} \quad (14)$$

To measure the ν_μ content of the beam, we introduce a “ ν_μ number” operator $\hat{\mu} \equiv |\mu\rangle\langle\mu|$. This operator acts only on the beam states and gives $\langle x|\hat{\mu}|x\rangle = \sin^2 \alpha$. To find the probability that the beam neutrino after interacting with N_2 background neutrinos will be measured as ν_μ , we compute the expectation value of $\hat{\mu}$ in the final state of Eq. (13)

$$\begin{aligned} \langle F|\hat{\mu}|F\rangle &= a^2 \langle x|\hat{\mu}|x\rangle \\ &\times (\langle exx...x| + \langle xex...x| + \dots) \\ &(|exx...x\rangle + |xex...x\rangle + \dots) \\ &= a^2 \sin^2 \alpha ((N_2^2 - N_2) \cos^2 \alpha + N_2). \end{aligned} \quad (15)$$

The last line is obtained by observing that in the sum there are N_2 “diagonal” terms of the type $\langle x...xex...x|x...xex...x\rangle = 1$ and $N_2^2 - N_2$ “off-diagonal” terms of the type $\langle x...xex...x|x...exx...x\rangle = \cos^2 \alpha$.

The result in Eq. (15) shows how the connection between a single scattering event and Eq. (8) is made. For a single scattering event, $N_2 = 1$, one indeed finds that

the conversion probability $P_{\nu_e \rightarrow \nu_\mu} \propto \sin^2 \alpha$ is maximized when the background consists of muon neutrinos. As the number of scattering events increases, however, the maximum of conversion efficiency shifts to values of $\alpha < \pi/2$. In the limit of large N_2 , one finds $P_{\nu_e \rightarrow \nu_\mu} \propto N_2^2 \sin^2 2\alpha$, precisely as predicted by Eq. (8).

We can now understand what happens to an electron neutrino propagating in a muon neutrino background. Eq. (15) shows that the conversion rate does not strictly vanish, as predicted by Eq. (8), but is only proportional to N_2 , not to N_2^2 . As discussed before, the proportionality to N_2 is a feature of incoherent scattering. The coherent scattering part is indeed strictly zero in this case: the states $|e\mu\mu\mu\dots\rangle$, $|\mu e\mu\mu\dots\rangle$, etc., are *mutually orthogonal*, and hence have no overlap that could be coherently amplified.

It can be readily seen that the part of the final state $|F\rangle$ that gets amplified contains the projection of the final background states onto the *initial* background state $|xxx\dots\rangle$. Decomposing the states in $|F_1\rangle$ according to

$$|x...xex\dots\rangle = \langle x|e\rangle|x...xxx\dots\rangle + \langle y|e\rangle|x...xyx\dots\rangle, \quad (16)$$

where y is the state orthogonal to x , $\langle y|x\rangle = 0$, we can write the result (15) as

$$\begin{aligned} \langle F|\hat{\mu}|F\rangle &= a^2 \langle x|\hat{\mu}|x\rangle \\ &\times (\langle e|x\rangle N_2 \langle xxx\dots x| + \\ &\langle e|y\rangle \langle yxx\dots x| + \langle e|y\rangle \langle xyx\dots x| + \dots) \\ &(\langle x|e\rangle N_2 |xxx\dots x\rangle + \\ &\langle y|e\rangle |yxx\dots x\rangle + \langle y|e\rangle |xyx\dots x\rangle + \dots) \\ &= a^2 \langle x|\hat{\mu}|x\rangle (N_2^2 |\langle x|e\rangle|^2 + N_2 |\langle y|e\rangle|^2). \end{aligned} \quad (17)$$

We now return to the problem of two intersecting beams, one containing electron neutrinos and another containing neutrinos in the flavor superposition state ν_x . Suppose N_1 neutrinos from the first beam interact with N_2 neutrinos from the second beam. As a result of the interaction, the system evolves over time δt according to

$$|eee\dots\rangle|xxx\dots\rangle \longrightarrow |F\rangle = |eee\dots\rangle|xxx\dots\rangle + ia|F_1\rangle, \quad (18)$$

where $|F_1\rangle$ contains a sum of terms with all possible exchanges,

$$\begin{aligned} |F_1\rangle &= (|xee\dots e\rangle + |exe\dots e\rangle + |eex\dots e\rangle + \dots) \\ &\times (|exx\dots x\rangle + |xex\dots x\rangle + |xex\dots x\rangle + \dots). \end{aligned} \quad (19)$$

As in the derivation of Eq. (17), to separate the coherent and incoherent parts, the states in $|F_1\rangle$ need to be projected on the corresponding initial states and orthogonal states. This is done using Eq. (16) and

$$|xee\dots\rangle = \langle e|x\rangle|eee\dots\rangle + \langle \mu|x\rangle|\mu ee\dots\rangle. \quad (20)$$

As a result, we obtain

$$\begin{aligned}
|F_1\rangle &= N_1 N_2 |\langle e|x\rangle|^2 |eee\dots\rangle |xxx\dots\rangle \\
&+ N_2 \langle \mu|x\rangle \langle x|e\rangle (|\mu ee\dots\rangle + |e\mu e\dots\rangle + \dots) |xxx\dots\rangle \\
&+ N_1 \langle e|x\rangle \langle y|e\rangle |eee\dots\rangle (|yxx\dots\rangle + |xyx\dots\rangle + \dots) \\
&+ \langle \mu|x\rangle \langle y|e\rangle (|\mu ee\dots\rangle + |e\mu e\dots\rangle + \dots) (|yxx\dots\rangle + |xyx\dots\rangle + \dots).
\end{aligned} \tag{21}$$

We observe that the term in the last line contains a sum of mutually orthogonal states. This term represents the part that will not amplify coherently, as can be seen by repeating the arguments given in connection with Eq. (17). We further note that if one *drops this term*, at first order

in *a* the final state equals

$$|F\rangle = |e'e'e'\dots e'e'\rangle |x'x'x'\dots x'x'\rangle, \tag{22}$$

where

$$|e'\rangle = |e\rangle + iN_2 a [1/2 \times |\langle x|e\rangle|^2 |e\rangle + \langle \mu|x\rangle \langle x|e\rangle |\mu\rangle], \tag{23}$$

$$|x'\rangle = |x\rangle + iN_1 a [1/2 \times |\langle e|x\rangle|^2 |x\rangle + \langle e|x\rangle \langle y|e\rangle |y\rangle]. \tag{24}$$

Eqs. (22,23,24) represent the central result of this section. They show that if we take the initial state to be a product of single-particle neutrino states and evolve it in time — carefully separating the coherent effects and dropping the incoherent ones — the final state obtained is again *a product of single particle states*, each rotated according to Eqs. (23,24). No coherent superposition of many-particle states is formed.

It is important to emphasize that to arrive at this conclusion we only needed to consider elementary scattering events (exchanges) between the beam and the background. No assumptions, such as decoherence between mass eigenstates in the background, were necessary.

IV. ONE-PARTICLE EVOLUTION EQUATIONS

A. Evolution equation: active-active oscillations

The construction in the preceding section can be used to obtain a differential equation describing the time evolution of a one-particle neutrino state. We first recall that in deriving Eq. (21) the flavor blind non-exchange interactions were omitted. It is easy to see that to include their effect one should add an additional term, $N_1 N_2 |eee\dots\rangle |xxx\dots\rangle$, to the final state $|F_1\rangle$ in Eq. (21),

$$\begin{aligned}
|F_1\rangle &= N_1 N_2 (1 + |\langle e|x\rangle|^2) |eee\dots\rangle |xxx\dots\rangle \\
&+ N_2 \langle \mu|x\rangle \langle x|e\rangle (|\mu ee\dots\rangle + |e\mu e\dots\rangle + \dots) |xxx\dots\rangle \\
&+ N_1 \langle e|x\rangle \langle y|e\rangle |eee\dots\rangle (|yxx\dots\rangle + |xyx\dots\rangle + \dots) \\
&+ \langle \mu|x\rangle \langle y|e\rangle (|\mu ee\dots\rangle + |e\mu e\dots\rangle + \dots) (|yxx\dots\rangle + |xyx\dots\rangle + \dots).
\end{aligned} \tag{25}$$

Eqs. (23) and (24), describing the evolution of the one-particle neutrino states over small time δt , then become

$$|e'\rangle = |e\rangle + iN_2 a [1/2 \times (1 + |\langle x|e\rangle|^2) |e\rangle + \langle \mu|x\rangle \langle x|e\rangle |\mu\rangle], \tag{26}$$

$$|x'\rangle = |x\rangle + iN_1 a [1/2 \times (1 + |\langle e|x\rangle|^2) |x\rangle + \langle e|x\rangle \langle y|e\rangle |y\rangle]. \tag{27}$$

We observe that, although Eq. (26) was obtained for the initial states ν_e and ν_x , the derivation used only par-

ticle exchanges and did not in any way rely on the par-

ticular choice of the initial states. Therefore, at any time t , if the initial state of the beam neutrino is ψ and the background neutrinos are all in the state ϕ , the evolution over a small time step is

$$|\psi(t + \delta t)\rangle - |\psi(t)\rangle = -i\sqrt{2}G_F N_2(1 - \cos\beta)/V\delta t \times [1/2 \times (1 + |\langle\phi|\psi\rangle|^2)|\psi\rangle + \langle\psi^\perp|\phi\rangle\langle\phi|\psi\rangle|\psi^\perp\rangle], \quad (28)$$

where $\psi^\perp$ is the state orthogonal to ψ and we restored the coefficients, including the angular factor.

Making use of the completeness relation $|\psi\rangle\langle\psi| + |\psi^\perp\rangle\langle\psi^\perp| = 1$, we rewrite Eq. (28) as

$$|\psi(t + \delta t)\rangle - |\psi(t)\rangle = -i\sqrt{2}G_F N_2(1 - \cos\beta)/V\delta t \times [1/2|\psi\rangle + |\phi\rangle\langle\phi|\psi\rangle - 1/2|\langle\phi|\psi\rangle|^2|\psi\rangle]. \quad (29)$$

The evolution of a one-particle neutrino state is therefore described by the following differential equation:

$$i\frac{d|\psi^{(i)}\rangle}{dt} = H_{aa}|\psi^{(i)}\rangle, \\ H_{aa} = \sum_j \frac{\sqrt{2}G_F}{V}(1 - \cos\beta^{(ij)}) \times \left[\frac{1}{2} + |\phi^{(j)}\rangle\langle\phi^{(j)}| - \frac{1}{2}|\langle\phi^{(j)}|\psi^{(i)}\rangle|^2 \right]. \quad (30)$$

Here index (i) refers to a given particle for which the equation is written, and (j) runs over all other neutrinos in the ensemble. The summation over the scattering angles $\beta^{(ij)}$ was introduced to make this equation applicable to a more general case of neutrinos propagating in different directions.

B. Evolution equation: active-sterile oscillations

The preceding analysis dealt with flavor conversions between active neutrino states. The method developed there can be extended to describe the conversions between active and sterile flavor states. In doing so, one should keep in mind two important differences between the two cases. In the active-sterile case: (i) only active components participate in the interactions, and (ii) the interaction amplitudes are proportional to the active content of the neutrino states involved. Performing calculations similar to those in Secs. III and IV A (see the Appendix) we find that the single-particle Hamiltonian for this case is given by

$$H_{as}^{(i)} = \sum_j \frac{2\sqrt{2}G_F}{V}(1 - \cos\beta^{(ij)}) \times |\langle\phi^{(j)}|e\rangle|^2[|e\rangle\langle e| - 1/2|\langle\psi^{(i)}|e\rangle|^2]. \quad (31)$$

As before, $\psi^{(i)}$ denotes the state of the neutrino for which the evolution equation is written and $\phi^{(j)}$ represents the flavor state of the j th neutrino in the background. It

is worth noting that this Hamiltonian includes both the effects of the t -channel and the u -channel diagrams.

For comparison, the standard Hamiltonian for an active-sterile neutrino system (the analogue of Eq. (3)) is

$$H_{as}^{(i)} = \sum_j \frac{2\sqrt{2}G_F}{V}(1 - \cos\beta^{(ij)}) \begin{pmatrix} \cos^2\alpha^{(j)} & 0 \\ 0 & 0 \end{pmatrix} \\ = \sum_j \frac{2\sqrt{2}G_F}{V}(1 - \cos\beta^{(ij)})|\langle\phi^{(j)}|e\rangle|^2|e\rangle\langle e|. \quad (32)$$

Here $\alpha^{(j)}$ is the mixing angle of the j th neutrino in the background, $\cos^2\alpha^{(j)} = |\langle\phi^{(j)}|e\rangle|^2$.

C. Background of antineutrinos

We now consider flavor transformation of the neutrino beam caused by the presence of antineutrinos in the background. For concreteness, let us envision a modification of the thought experiment depicted in Fig. 2 in which the beam contains neutrinos in a superposition of flavor states, $\nu_z = \cos\beta\nu_e + \sin\beta\nu_\mu$, and the background contains *antineutrinos* in a flavor superposition $\bar{\nu}_x = \cos\alpha\bar{\nu}_e + \sin\alpha\bar{\nu}_\mu$. To fix the notation, let ν_w be the flavor state orthogonal to ν_z , $\langle\nu_w|\nu_z\rangle = 0$ and $\bar{\nu}_y$ be the flavor state orthogonal to $\bar{\nu}_x$, $\langle\bar{\nu}_y|\bar{\nu}_x\rangle = 0$.

Consider what elementary processes are possible in this case. As can be easily seen, in addition to the t - and u -channel processes, $\nu_z\bar{\nu}_x \rightarrow \nu_z\bar{\nu}_x$ and $\nu_z\bar{\nu}_x \rightarrow \bar{\nu}_x\nu_z$, it is possible to have s -channel annihilation diagrams $\nu_z\bar{\nu}_z \rightarrow \nu_z\bar{\nu}_z, \nu_w\bar{\nu}_w, \bar{\nu}_z\nu_z, \bar{\nu}_w\nu_w$. Notice that only the $\bar{\nu}_z$ component of the $\bar{\nu}_x$ state participates in the s -channel processes.

The t -channel process is flavor-diagonal and hence does not cause flavor conversion, just like the corresponding process in the neutrino background. The u -channel process puts an antineutrino in the beam and, therefore, cannot interfere coherently with the incident neutrino wave. Any coherent flavor changes, therefore, can only be due to the s -channel process.

Since only the $\bar{\nu}_z$ component of the $\bar{\nu}_x$ state participates in the s -channel process, the beam neutrinos will not change flavor if the background antineutrinos are in the orthogonal flavor state, $\bar{\nu}_w$. While this is similar to what was found for a background of neutrinos in the state ν_w , on the microscopical level the two cases are quite different. For the ν_w background, the amplitude of flavor conversion for a single elementary event is nonzero, but the conversion rate is only proportional to N , because the amplitudes add up incoherently. By comparison, for the $\bar{\nu}_w$ background already the elementary amplitude vanishes and the ν_w appearance rate in the beam is strictly zero. (Instead, $\bar{\nu}_w$ *antineutrinos* will appear in the beam at the rate proportional to N , due to the u -channel process.)

The elementary scattering event can be written as

$$|\nu_z \bar{\nu}_x\rangle \longrightarrow |\nu_z \bar{\nu}_x\rangle - ia\langle \bar{\nu}_z | \bar{\nu}_x \rangle (|\nu_z \bar{\nu}_z\rangle + |\nu_w \bar{\nu}_w\rangle), \quad (33)$$

where the t -channel process as well as the processes that put an antineutrino in the beam have been omitted. The minus sign appears because the amplitudes for neutrino-neutrino and neutrino-antineutrino scattering processes have opposite signs.

As before, we project the final state on the initial states and orthogonal states,

$$\begin{aligned} |\nu_z \bar{\nu}_x\rangle \longrightarrow & |\nu_z \bar{\nu}_x\rangle - ia\langle \bar{\nu}_z | \bar{\nu}_x \rangle \times \\ & (\langle \bar{\nu}_x | \bar{\nu}_z \rangle |\nu_z \bar{\nu}_x\rangle + \langle \bar{\nu}_y | \bar{\nu}_z \rangle |\nu_z \bar{\nu}_y\rangle + \\ & \langle \bar{\nu}_x | \bar{\nu}_w \rangle |\nu_w \bar{\nu}_x\rangle + \langle \bar{\nu}_y | \bar{\nu}_w \rangle |\nu_w \bar{\nu}_y\rangle). \end{aligned} \quad (34)$$

The rest of the argument proceeds in complete analogy to the case of the neutrino background. The first three terms in parentheses in Eq. (34) will amplify coherently and the last term will not. Summing over many elementary scattering events we obtain an expression similar to Eq. (21), which gives a one-particle evolution equation

$$\begin{aligned} i \frac{d|\psi^{(i)}\rangle}{dt} &= H_{a\bar{a}} |\psi^{(i)}\rangle, \\ H_{a\bar{a}} &= - \sum_j \frac{\sqrt{2}G_F}{V} (1 - \cos \beta^{(ij)}) \times \\ &\quad \left[\frac{1}{2} + |\bar{\phi}^{(j)}\rangle \langle \bar{\phi}^{(j)}| - \frac{1}{2} |\langle \bar{\phi}^{(j)} | \psi^{(i)} \rangle|^2 \right], \end{aligned} \quad (35)$$

where $\bar{\phi}^{(j)}$ denotes the flavor state of the j th antineutrino in the background and the contributions of the t -channel processes have been included. Thus, the effects of the neutrino and antineutrino backgrounds on the coherent neutrino flavor evolution have exactly the same form (but opposite signs!), even though at the microscopical level the two cases are different.

When both neutrinos and antineutrinos are present in the background, their refractive effects add up linearly. Therefore, the Hamiltonian describing the flavor evolution of a neutrino beam equals the sum of the two contributions, Eqs. (30) and (35). Further generalization to include the effects of other matter (electrons, nucleons) and vacuum oscillations is straightforward. Just like in the case of the usual MSW effect, one should add the Hamiltonian terms H_{vac} and H_{mat} to the neutrino induced Hamiltonian (see Eq. (1)).

D. Comparison to the standard results

We now compare the one-particle Hamiltonians we have obtained, Eqs. (30), (31), and (35), to the corresponding accepted results, Eqs. (3) and (32). We see that, while the accepted results are similar to ours, there are important differences: in all three cases, our Hamiltonians contain additional terms. It is important to understand both the origin of this difference and whether it leads to any physical consequences.

First, we would like to establish whether the presence of the additional terms in our results affects the flavor evolution of the neutrino system. As can be readily seen, in all three cases the additional terms are proportional to the identity matrix in the flavor space. The evolution equations thus have the form

$$i\psi' = (H_0 + C(\psi, \phi)\mathbb{I})\psi, \quad (36)$$

where H_0 is the “standard” Hamiltonian given in Eq. (3) or (32) and $\mathbb{I}$ is the identity matrix. We observe that, if $\psi_0(t)$ is the solution of the equation $i\psi' = H_0\psi$, then $\psi_1(t)$ given by

$$\psi_1(t) = \exp \left[-i \int^t C(\psi_0(\tilde{t}), \phi_0(\tilde{t})) d\tilde{t} \right] \psi_0(t) \quad (37)$$

is a solution of Eq. (36). This means that the extra terms give an overall shift to both energy levels, without affecting neutrino flavor evolution.

Indeed, the physical origin of the difference in all three cases can be traced to the part of the interaction that *does not change flavor*. For concreteness, we for a moment specialize to the first of the three cases, the active-active conversions in the neutrino background. It proves instructive to return to the evolution of the beam neutrino over an infinitesimal time step. In our case, the result is given by Eq. (26), while the accepted result, Eq. (3), gives

$$|e'\rangle = |e\rangle + iN_2a[(1 + \cos^2 \alpha)|e\rangle + \sin \alpha \cos \alpha |\mu\rangle]. \quad (38)$$

The difference between the two is the factor of $1/2$, which multiplies the state $|e\rangle$ (the flavor-preserving part) in the brackets in Eq. (26). This factor, in turn, can be traced to Eq. (25): it comes from the first term in $|F_1\rangle$, which *must be split* between the beam and the background to avoid overcounting. This is the origin of the factors of $1/2$ in Eqs. (26) and (27).

The situation is not unlike what happens in electrostatics. The interaction energy in a system of charges is given by $1/2 \sum_i q_i \phi_i$ and the factor of $1/2$ ensures that the interaction energy between pairs of charges is not counted twice. In our case, the extra terms serve the same purpose, to prevent counting the interaction energy twice. This can be seen as follows. Both active-active and active-sterile evolution equations are particular cases of a general case when the two flavor states have different couplings to the Z boson. As shown in the Appendix, the evolution equation in this general case can be written in a form (57):

$$i \frac{d|\psi^{(i)}\rangle}{dt} = \left[H_0^{(i)} - \frac{1}{2} \langle \psi^{(i)} | H_0^{(i)} | \psi^{(i)} \rangle \right] |\psi^{(i)}\rangle, \quad (39)$$

where $H_0^{(i)}$ is the generalization of the standard Hamiltonian [10],

$$H_0^{(i)} = \sum_j \frac{\sqrt{2}G_F}{V} (1 - \cos \beta^{(ij)}) \left[G(\eta) |\phi^{(j)}\rangle \langle \phi^{(j)}| G(\eta) + G(\eta) \langle \phi^{(j)}| G(\eta) |\phi^{(j)}\rangle \right], \quad (40)$$

with G being the matrix of couplings,

$$G(\eta) = \begin{pmatrix} 1 & 0 \\ 0 & \eta \end{pmatrix}. \quad (41)$$

The second term in the evolution equation (39) has the form of the expectation value $1/2 \langle \psi^{(i)} | H_0^{(i)} | \psi^{(i)} \rangle$. This form makes explicit the physical meaning of this term as a correction to avoid double counting of the energy of the system. It is worth emphasizing that this term is *always* proportional to the identity in the flavor space, even when the two flavor states have different couplings. Physically, this is because the part of the interaction that needs to be split is always the part that conserves flavor (see the Appendix).

The extra term, $1/2 \langle \psi^{(i)} | H_0^{(i)} | \psi^{(i)} \rangle$, depends not only on the state of the background but also on the state of the beam itself. We therefore caution the reader that the superposition principle, which is always used for the MSW effect in normal matter, does not apply in this case. For instance, suppose that a beam neutrino which is ν_e at $t = 0$ as a result of the evolution becomes a state ν' and, similarly, a beam neutrino which is ν_μ at $t = 0$ becomes a state ν'' . Then, it is in general not true that the state $\nu_x = \cos \alpha \nu_e + \sin \alpha \nu_\mu$ will become $\cos \alpha \nu' + \sin \alpha \nu''$.

It would be incorrect to conclude that the extra terms have no physical effect whatsoever. While they indeed do not change neutrino flavor evolution, they do modify the absolute value of the refraction index of a neutrino medium and, hence, at least in principle, change the bending of a neutrino beam in a dense neutrino medium with a density gradient. This effect is present even if there is only one neutrino flavor in the system.

V. MORE ON THE ENTANGLED SYSTEM

We have shown that if the neutrino system initially does not contain entangled states, such states are not formed as a result of coherent evolution in the system. It can be argued, moreover, that such evolution can lead to an effective loss of coherence between entangled states. To illustrate this, let us consider a beam in an entangled flavor state,

$$|\text{ent}\rangle \equiv (|xxx\dots\rangle + |yyy\dots\rangle)/\sqrt{2}, \quad (42)$$

(here x and y are not necessarily orthogonal) propagating in the (unentangled) background $|zzz\dots\rangle$.

At time $t = 0$ the expectation value of some operator, for example, the ν_μ number operator, $\hat{\mu} = \sum |\mu\rangle \langle \mu|$,

where the sum runs over all particles in the beam, is given by

$$\begin{aligned} \langle \text{ent} | \hat{\mu} | \text{ent} \rangle &= \frac{1}{2} (\langle xxx\dots | \hat{\mu} | xxx\dots \rangle + \langle yyy\dots | \hat{\mu} | yyy\dots \rangle \\ &+ \langle xxx\dots | \hat{\mu} | yyy\dots \rangle + \langle yyy\dots | \hat{\mu} | xxx\dots \rangle). \end{aligned} \quad (43)$$

The first two terms on the right hand side simply count the muon neutrino content in the states $|xxx\dots\rangle$ and $|yyy\dots\rangle$ and the last two terms represent the effect of entanglement.

Let us consider the effects of time evolution on the expectation value of $\hat{\mu}$. Each of the two terms in $|\text{ent}\rangle$ is a product of single-particle states and according to our earlier findings over time will remain a product of single-particle states. Let us write the state at time $t = t_1$ as

$$(|x'x'x'\dots\rangle |z'z'z'\dots\rangle + |y''y''y''\dots\rangle |z''z''z''\dots\rangle)/\sqrt{2}, \quad (44)$$

where z' , z'' , x' , and y'' are the results of solving a system of equations given in Eq. (30). (For example, z' and x' are found by solving the equations for the initial state $|xxx\dots\rangle |zzz\dots\rangle$.)

The expectation value of $\hat{\mu}$ in the state (44) is given by

$$\begin{aligned} &\frac{1}{2} (\langle x'x'x'\dots | \hat{\mu} | x'x'x'\dots \rangle + \langle y''y''y''\dots | \hat{\mu} | y''y''y''\dots \rangle \\ &+ \langle x'x'x'\dots | \hat{\mu} | y''y''y''\dots \rangle \langle z'z'z'\dots | z''z''z''\dots \rangle \\ &+ \langle y''y''y''\dots | \hat{\mu} | x'x'x'\dots \rangle \langle z''z''z''\dots | z'z'z'\dots \rangle). \end{aligned} \quad (45)$$

Since the states z' and z'' will generically be different, the absolute value of the inner product $\langle z'' | z' \rangle$ will be < 1 . The last two terms in Eq. (45) therefore contain suppression factors $|\langle z'' | z' \rangle|^{N_2}$ and vanish as the number of neutrinos in the background N_2 is taken to infinity. As already mentioned, these terms represent the entanglement between the two states; the system therefore behaves as if the beam was an incoherent mixture of $|x'x'x'\dots\rangle$ and $|y''y''y''\dots\rangle$.

Of course, rigorously speaking, the entanglement information is not completely lost in the system. It may happen that at some time t the states z' and z'' will be such that $|\langle z'' | z' \rangle| = 1$. In this case, the entanglement effect will reappear. We, however, regard this as an artificial arrangement and therefore maintain that for practical purposes the coherence is destroyed.

It is curious to note that, as the entanglement effect reappears, the phases of the states $|x'x'x'\dots\rangle |z'z'z'\dots\rangle$ and $|y''y''y''\dots\rangle |z''z''z''\dots\rangle$ will have an effect on the expectation value of $\hat{\mu}$ and hence the phases due to the additional term, introduced in Eq. (30) to avoid overcounting of energies, will have a physical effect.

VI. CONCLUSIONS

In summary, we have developed a conceptually simple, physical picture of coherent neutrino evolution in a medium of neutrinos and have shown how coherent interference of many elementary scattering events gives rise to the refraction phenomenon. Unlike the case of ordinary matter, in which coherent scattering leaves the background unchanged, in the neutrino background the scattering events change the state of the background, thus requiring a different type of analysis. We have found that only part of the elementary scattering amplitude is amplified coherently. This explains certain seemingly paradoxical results, such as why a neutrino traveling through the medium of neutrinos of opposite flavor does not undergo coherent flavor conversion.

We have shown that refraction does not lead to the creation of entangled states in a neutrino system, *i.e.*, if the state of the system is initially described by a product of single-particle states, the state remains a product of single-particle states as the system is evolved in time. Furthermore, the evolution effectively destroys initial entanglement in the system. It follows that for each neutrino the result of the coherent evolution can be described by a one-particle Schrödinger equation, as was assumed *a priori* in the literature.

We have derived the one-particle equation for active-active and active-sterile flavor transformation scenarios for a neutrino in a neutrino background. We also derived the equation for a neutrino in an antineutrino background. In all these cases, we found that in order to avoid overcounting of the interaction energy one has to introduce an extra term in the evolution equations that is not

present in the standard analyses. We have proven that this extra term does not affect the flavor evolution under normal conditions. It does, however, affect the value of the refraction index and hence the bending of a neutrino beam in a dense neutrino medium.

Acknowledgments

We thank Georg Raffelt, James Pantaleone, and John Bahcall for stimulating discussions and for valuable comments on the manuscript. We also acknowledge valuable comments from John Cornwall, Wick Haxton, Maxim Perelstein, Raymond Sawyer, and Raymond Volkas. A. F. is a Feynman Fellow at LANL. A. F. was supported at LANL by the Department of Energy, under contract W-7405-ENG-36, and at IAS by the Keck Foundation. C. L. was supported by the Keck Foundation and by the National Science Foundation grant PHY-0070928. We acknowledge the hospitality of the Kavli Institute for Theoretical Physics, where this research was supported in part by the National Science Foundation under Grant No. PHY99-07949.

Appendix: the cases of active-sterile neutrinos and neutrinos with different couplings

Let us derive the result (31), which refers to a system of one active and one sterile neutrino, ν_e and ν_s . We start by considering the elementary interaction between two neutrinos, as discussed in Sec. II B. The analogue of Eq. (9) in this case is

$$i \frac{d}{dt} \begin{pmatrix} |\nu_e(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_e(\vec{k})\nu_s(\vec{p})\rangle \\ |\nu_s(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_s(\vec{k})\nu_s(\vec{p})\rangle \end{pmatrix} = \frac{\sqrt{2}G_F}{V}(1 - \cos\beta) \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} |\nu_e(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_e(\vec{k})\nu_s(\vec{p})\rangle \\ |\nu_s(\vec{k})\nu_e(\vec{p})\rangle \\ |\nu_s(\vec{k})\nu_s(\vec{p})\rangle \end{pmatrix}, \quad (46)$$

which shows that, given an initial state $|\nu_z(\vec{k})\nu_x(\vec{p})\rangle$, only its active-active component, $|\nu_e(\vec{k})\nu_e(\vec{p})\rangle$, is affected by the evolution:

$$|\nu_z(\vec{k})\nu_x(\vec{p})\rangle \Rightarrow |\nu_z(\vec{k})\nu_x(\vec{p})\rangle - i dt \frac{2\sqrt{2}G_F}{V}(1 - \cos\beta) \langle e|z\rangle \langle e|x\rangle |\nu_e(\vec{k})\nu_e(\vec{p})\rangle. \quad (47)$$

Next, we apply this result to the case of two orthogonal neutrino beams, in the spirit of what was done in Sec. III. We take neutrinos in the first and second beams to be in the states $|z\rangle$ and $|x\rangle$, respectively, and omit neutrino momenta for simplicity. Similarly to Eq. (18), we get

$$|zzz\dots\rangle|xxx\dots\rangle \Rightarrow |F\rangle = |zzz\dots\rangle|xxx\dots\rangle + i2a|F_1\rangle, \quad (48)$$

where

$$\begin{aligned} |F_1\rangle &= \langle e|z\rangle \langle e|x\rangle (|ezz\dots z\rangle + |zez\dots z\rangle + |zze\dots z\rangle + \dots) \\ &\times (|exx\dots x\rangle + |xex\dots x\rangle + |xxe\dots x\rangle + \dots). \end{aligned} \quad (49)$$

Notice that the effects of both the u -channel and t -channel diagrams are included above.

One can then follow the same procedure as in Sec. III (see Eq. (21)) and decompose $|F_1\rangle$ as follows:

$$\begin{aligned}
|F_1\rangle = & \langle e|z\rangle\langle e|x\rangle[N_1N_2\langle z|e\rangle\langle x|e\rangle|zzz\dots\rangle|xxx\dots\rangle \\
& + N_2\langle w|e\rangle\langle x|e\rangle(|wzz\dots\rangle + |zwz\dots\rangle + \dots)|xxx\dots\rangle \\
& + N_1\langle z|e\rangle\langle y|e\rangle|zzz\dots\rangle(|yxx\dots\rangle + |xyx\dots\rangle + \dots) \\
& + \langle w|e\rangle\langle y|e\rangle(|wzz\dots\rangle + |zwz\dots\rangle + \dots)(|yxx\dots\rangle + |xyx\dots\rangle + \dots)
\end{aligned} \tag{50}$$

where $|y\rangle$ and $|w\rangle$ are the orthogonal states to $|x\rangle$ and $|z\rangle$ respectively ($\langle x|y\rangle = \langle z|w\rangle = 0$). The last term in Eq. (50) is not coherently enhanced and therefore can be dropped. This allows us to obtain, at first order in a , a factorized form:

$$|F\rangle \simeq |z'z'z'\dots\rangle|x'x'x'\dots\rangle, \tag{51}$$

where

$$\begin{aligned}
|z'\rangle &= |z\rangle + iN_2a|\langle e|z\rangle|^2|\langle e|x\rangle|^2|z\rangle + i2N_2a|\langle e|x\rangle|^2\langle w|e\rangle\langle e|z\rangle|w\rangle \\
&= |z\rangle + i2N_2a\langle e|z\rangle|\langle e|x\rangle|^2|e\rangle - iN_2a|\langle e|z\rangle|^2|\langle e|x\rangle|^2|z\rangle \\
|x'\rangle &= |x\rangle + iN_1a|\langle e|z\rangle|^2|\langle e|x\rangle|^2|x\rangle + i2N_1a|\langle e|z\rangle|^2\langle y|e\rangle\langle e|x\rangle|y\rangle \\
&= |x\rangle + i2N_1a\langle e|x\rangle|\langle e|z\rangle|^2|e\rangle - iN_1a|\langle e|z\rangle|^2|\langle e|x\rangle|^2|x\rangle.
\end{aligned} \tag{52}$$

From this a one-particle equation follows:

$$\begin{aligned}
i\frac{d|\psi^{(i)}\rangle}{dt} &= \sum_j \frac{2\sqrt{2}G_F}{V}(1 - \cos\beta^{(ij)}) \\
&\times |\langle\phi^{(j)}|e\rangle|^2 \left[|e\rangle\langle e| - 1/2|\langle\psi^{(i)}|e\rangle|^2 \right] |\psi^{(i)}\rangle,
\end{aligned} \tag{53}$$

which proves the result (31).

We now discuss the generalization of our findings to two active neutrinos with different couplings to the Z boson. The study of this case provides a unified description of the results for active-active and active-sterile cases we have discussed so far. Furthermore, it allows us to compare our results with the corresponding discussion given

in ref. [10].

Let us consider two neutrino states, ν_e and ν_ρ , and take ν_e as having the ordinary Standard Model coupling, g_e , with the Z boson. The coupling of ν_ρ with the Z is defined as $g_\rho \equiv \eta g_e$. In terms of η , the two-neutrinos equation, (46), is generalized by the replacements $\nu_s \rightarrow \nu_\rho$ and

$$\begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow B(\eta) \equiv \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & \eta & \eta & 0 \\ 0 & \eta & \eta & 0 \\ 0 & 0 & 0 & 2\eta^2 \end{pmatrix}. \tag{54}$$

It is manifest that the active-active (Eq. (9)) and active-sterile (Eq. (46)) cases are recovered for $\eta = 1$ and $\eta = 0$ respectively. The generalization of Eq. (50) has the form

$$\begin{aligned}
|F_1\rangle = & N_1N_2\langle x|\langle z|B(\eta)|z\rangle|x\rangle|zzz\dots\rangle|xxx\dots\rangle \\
& + N_2\langle x|\langle w|B(\eta)|z\rangle|x\rangle(|wzz\dots\rangle + |zwz\dots\rangle + \dots)|xxx\dots\rangle \\
& + N_1\langle y|\langle z|B(\eta)|z\rangle|x\rangle|zzz\dots\rangle(|yxx\dots\rangle + |xyx\dots\rangle + \dots) \\
& + \langle y|\langle w|B(\eta)|z\rangle|x\rangle(|wzz\dots\rangle + |zwz\dots\rangle + \dots)(|yxx\dots\rangle + |xyx\dots\rangle + \dots).
\end{aligned} \tag{55}$$

Once the last term in Eq. (55) is neglected, as discussed in Sec. III, one gets the effective one-particle equations:

$$\begin{aligned}
|z'\rangle &= |z\rangle + \frac{1}{2}iN_2a\langle x|\langle z|B(\eta)|z\rangle|x\rangle|z\rangle + iN_2a\langle x|\langle w|B(\eta)|z\rangle|x\rangle|w\rangle \\
|x'\rangle &= |x\rangle + \frac{1}{2}iN_1a\langle x|\langle z|B(\eta)|z\rangle|x\rangle|x\rangle + iN_1a\langle y|\langle z|B(\eta)|z\rangle|x\rangle|y\rangle.
\end{aligned} \tag{56}$$

Notice that, as before, the factors of $1/2$ arise as a result of splitting of the first (flavor conserving) term in

Eq. (55).

Using the form (54), and working out the matrix elements in Eq. (56), it is possible to obtain the generalization of the result (53). We get

$$i \frac{d|\psi^{(i)}\rangle}{dt} = \left[H_0^{(i)} - \frac{1}{2} \langle \psi^{(i)} | H_0^{(i)} | \psi^{(i)} \rangle \right] |\psi^{(i)}\rangle, \quad (57)$$

$$H_0^{(i)} = \sum_j \frac{\sqrt{2}G_F}{V} (1 - \cos \beta^{(ij)}) \left[G(\eta) |\phi^{(j)}\rangle \langle \phi^{(j)}| G(\eta) + G(\eta) \langle \phi^{(j)}| G(\eta) |\phi^{(j)}\rangle \right]. \quad (58)$$

Here the angular factors $(1 - \cos \beta^{(ij)})$ have been restored to account for general orientations of the neutrino momenta, and we use the matrix of the couplings (normalized to g_e) in the (ν_e, ν_ρ) basis:

$$G(\eta) = \begin{pmatrix} 1 & 0 \\ 0 & \eta \end{pmatrix}. \quad (59)$$

Our result (58) coincides with the one-particle Hamiltonian given in ref. [10].

As an example, we give the explicit expression of $H_0^{(i)}$ for the case of orthogonal beams with the neutrinos in the

where $H_0^{(i)}$ has the form

second beam being all in the same state, $\phi^{(j)} = \cos \alpha \nu_e + \sin \alpha \nu_\rho$:

$$H_0^{(i)} = \frac{\sqrt{2}G_F N_2}{V} \times \begin{pmatrix} 2 \cos^2 \alpha + \eta \sin^2 \alpha & \eta \cos \alpha \sin \alpha \\ \eta \cos \alpha \sin \alpha & \eta \cos^2 \alpha + 2 \eta^2 \sin^2 \alpha \end{pmatrix}. \quad (60)$$

It is easy to see that Eq. (60) reproduces the limiting cases of active-active ($\eta = 1$) and active-sterile ($\eta = 0$) neutrinos, Eqs. (7) and (32).

-
- [1] A. Y. Smirnov and S. P. Mikheev (1986), in *Proceedings of the Sixth Moriond Workshop on Massive Neutrinos in Astrophysics and in Particle Physics*, Tignes, France, 1986, pp. 355-372.
 - [2] G. M. Fuller, R. W. Mayle, J. R. Wilson, and D. N. Schramm, *Astrophys. J.* **322**, 795 (1987).
 - [3] D. Notzold and G. Raffelt, *Nucl. Phys.* **B307**, 924 (1988).
 - [4] R. Barbieri and A. Dolgov, *Phys. Lett.* **B237**, 440 (1990).
 - [5] M. J. Savage, R. A. Malaney, and G. M. Fuller, *Astrophys. J.* **368**, 1 (1991).
 - [6] R. Barbieri and A. Dolgov, *Nucl. Phys.* **B349**, 743 (1991).
 - [7] P. Langacker and J. Liu, *Phys. Rev.* **D46**, 4140 (1992), hep-ph/9206209.
 - [8] J. Pantaleone, *Phys. Rev.* **D46**, 510 (1992).
 - [9] J. Pantaleone, *Phys. Lett.* **B287**, 128 (1992).
 - [10] G. Sigl and G. Raffelt, *Nucl. Phys.* **B406**, 423 (1993).
 - [11] B. H. J. McKellar and M. J. Thomson, *Phys. Rev.* **D49**, 2710 (1994).
 - [12] V. A. Kosteletsky, J. Pantaleone, and S. Samuel, *Phys. Lett.* **B315**, 46 (1993).
 - [13] V. A. Kosteletsky and S. Samuel, *Phys. Lett.* **B318**, 127 (1993).
 - [14] V. A. Kosteletsky and S. Samuel, *Phys. Rev.* **D49**, 1740 (1994).
 - [15] V. A. Kosteletsky and S. Samuel, *Phys. Rev.* **D52**, 3184 (1995), hep-ph/9507427.
 - [16] C. Lunardini and A. Y. Smirnov, *Phys. Rev.* **D64**, 073006 (2001), hep-ph/0012056.
 - [17] S. Hannestad, *Phys. Rev.* **D65**, 083006 (2002), astro-ph/0111423.
 - [18] A. D. Dolgov et al., *Nucl. Phys.* **B632**, 363 (2002), hep-ph/0201287.
 - [19] Y. Y. Y. Wong, *Phys. Rev.* **D66**, 025015 (2002), hep-ph/0203180.
 - [20] K. N. Abazajian, J. F. Beacom, and N. F. Bell, *Phys. Rev.* **D66**, 013008 (2002), astro-ph/0203442.
 - [21] J. Pantaleone, *Phys. Lett.* **B342**, 250 (1995), astro-ph/9405008.
 - [22] Y. Z. Qian and G. M. Fuller, *Phys. Rev.* **D51**, 1479 (1995), astro-ph/9406073.
 - [23] G. Sigl, *Phys. Rev.* **D51**, 4035 (1995), astro-ph/9410094.
 - [24] G. C. McLaughlin, J. M. Fetter, A. B. Balantekin, and G. M. Fuller, *Phys. Rev.* **C59**, 2873 (1999), astro-ph/9902106.
 - [25] S. Pastor and G. Raffelt, *Phys. Rev. Lett.* **89**, 191101 (2002), astro-ph/0207281.
 - [26] S. Samuel, *Phys. Rev.* **D48**, 1462 (1993).
 - [27] S. Samuel, *Phys. Rev.* **D53**, 5382 (1996), hep-ph/9604341.
 - [28] J. Pantaleone, *Phys. Rev.* **D58**, 073002 (1998).
 - [29] S. Pastor, G. G. Raffelt, and D. V. Semikoz, *Phys. Rev.* **D65**, 053011 (2002), hep-ph/0109035.
 - [30] For physical applications, both in the case of the early Universe and supernova, the relevant neutrino energies are in the $10^6 - 10^7$ eV range.
 - [31] To avoid conflict with the Pauli's principle and to make possible labelling of neutrinos by their momenta, we can assume that neutrinos have slightly different energies.